



SAM Contributes to SASSI-3 Accuracy

This SASSI-3 profile of a 37-year-old female was called in to our clinical support line. As we look at her results, it appears that she answered the items in a meaningful manner (RAP=0) and there is no evidence of defensive responding (DEF=3). She is likely to have a substance dependence disorder based on Rule 7 (FVOD=15, SAM=8).

Notice that despite the relatively low DEF score and apparent lack of defensive responding, the SAM scale score, when combined with the elevated FVOD score, leads to a test positive result. While it is true that the SAM scale score plays a vital role in the accuracy of the decision rules in this case, it is important to remember that the clinical meaning of this score is unclear. Therefore, it would be inappropriate to draw any clinical inference from the fact that the SAM score is elevated.

The client acknowledges significant problems related to her use of drugs other than alcohol. She is likely to have experienced some loss of control, negative consequences, and increased tolerance as a result of her substance misuse. However, her average OAT score (OAT=5) may be an indication of some limited ability or willingness to acknowledge behavioral problems commonly associated with substance dependence.

The moderately elevated SYM (SYM=6) is consistent with clients who often are not able to recognize the manner in which substance dependence is manifested in their lives. Her responses are similar to individuals who live in a social milieu where substance abuse and its related consequences are fairly routine and normalized. This experience may limit her ability to characterize her substance usage as problematic. Indeed, she may be somewhat surprised that the SASSI results could even indicate an addiction problem.

She may also have poor insight and self-awareness regarding the way in which substance dependence has pervaded into numerous areas of her life (SAT=5, SAT greater than OAT).

She may need assistance in recognizing the ways in which her substance use has effected her psychological, emotional and spiritual functioning. Her scores are consistent with individuals who tend to present as relatively well functioning, maintaining an illusion of control over their thoughts, feelings and actions.

The client's responses are similar to those of individuals who are experiencing emotional pain (DEF=3). Individuals who score in this range tend to be overly self-

critical, may experience depressive symptoms and sometimes report a history of trauma. She may be quite limited in her ability to recognize personal strengths, focusing more on limitations, failures and feelings of low self-worth.

Treatment Recommendations:

This client is likely to have a substance dependence disorder and should be considered for relatively intensive addictions treatment. A comprehensive behavioral health evaluation may be necessary to rule out the need for additional psychiatric intervention. Although she demonstrates some ability to acknowledge relevant behavioral symptoms of her addiction, a viable treatment plan should include initial efforts to increase her self-awareness and insight into the full nature of her substance dependence problems. Education and other cognitively based interventions may be helpful.

Most likely, she will need help in recognizing that her misuse of alcohol and other drugs is similar to that of other substance dependent people. A content analysis of her responses on the FVOD and SYM items may be one way to help her realize that it is in her best interest and within her capacity to change. Interventions geared towards joining her in the process of getting in touch with her emotional pain and how it relates to her substance dependency would be important to consider,


Community-based self-help support groups could provide additional encouragement and support.

SASSI-3 Substance Abuse Subtle Screening Inventory

For free consultation on this profile 1-888 BY SASSI 1-888-297-2774 M-Th 8-6 Fri 8-5 EST



Name _____ Gender **F** Age **37**
 Case Number _____ Test date _____



RAP Random Answering Pattern

0 — Check if RAP is 2 or more. Results may not be meaningful. Try to resolve problem before proceeding. Such profiles have an average decreased accuracy of approximately 20%.

Check every rule, yes or no.

Rule 1 FVA 20 or more? yes no

Rule 2 FVOD 21 or more? yes no

Rule 3 SYM 7 or more? yes no

Rule 4 OAT 10 or more? yes no

Rule 5 SAT 6 or more? yes no

Rule 6 OAT 7 or more and SAT 5 or more Both? yes no

Rule 7 FVA 9 or more OR FVOD 15 or more and SAM 8 or more Both? yes no

Rule 8 OAT 5 or more and DEF 8 or more and SAM 8 or more All three? yes no

Rule 9 FVA 14 or more OR FVOD 8 or more and SAT 2 or more and DEF 4 or more and SAM 4 or more All four? yes no

THE DECISION RULE:

ANY rule answered "yes"?

HIGH PROBABILITY of having a Substance Dependence Disorder

ALL rules answered "no"?

LOW PROBABILITY of having a Substance Dependence Disorder

Check if DEF is 8 or more. Elevated DEF scores increase the possibility of the SASSI missing substance dependent individuals. Elevated DEF may also reflect situational factors.

Copyright © 1994, 1997 Glenn A. Miller

v 97F 3/26/97